


Campylobacteriosis Incidence Snohomish County, 2002 - 2008

- From 2002 through 2008, the incidence of campylobacteriosis in both Snohomish County and Washington State has not significantly changed.
- In 2008 the Snohomish County rate was comparable with the Washington State rate.



Year	Snohomish County				Washington State		
	Cases	Population	Cases per 100,000	95% C.I. ¹	Cases	Population	Cases per 100,000
2002	105	627,974	16.7	(13.7, 20.2)	1,024	6,041,710	17.0
2003	96	637,500	15.1	(12.2, 18.4)	929	6,098,300	15.2
2004	88	644,811	13.7	(11.0, 16.8)	847	6,167,800	13.7
2005	110	655,800	16.8	(13.8, 20.2)	1,036	6,256,400	16.6
2006	94	671,836	14.0	(11.3, 17.1)	983	6,375,600	15.4
2007	117	686,300	17.1	(14.1, 20.4)	1,020	6,488,000	15.7
2008	123	696,629	17.7	(14.7, 21.1)	1,069	6,587,600	16.2

¹ 95% Confidence Interval: A confidence interval is a range of values that has a specified probability of containing the rate. Narrow confidence intervals indicate that rate is precisely estimated, while wider confidence intervals indicate that the estimated rate is less precisely estimated.