


## Smoking During Pregnancy – Snohomish County, 1999 - 2008

- In 2008 the rates of women smoking during pregnancy were comparable between Snohomish County women and Washington State women.
- The percent of women smoking during pregnancy significantly declined between 1999 and 2004 in both Snohomish County and Washington State.



Year	Snohomish County				Washington State		
	# of Pregnant Women Smoking	Pregnant Women	% of Pregnant Women Smoking	95% C.I. <sup>1</sup>	# of Pregnant Women Smoking	Pregnant Women	% of Pregnant Women Smoking
1999	1,063	8,046	13.2	(12.4, 14.0)	10,586	74,410	14.2
2000	997	8,251	12.1	(11.3, 12.9)	10,450	77,384	13.5
2001	1,020	8,483	12.0	(11.3, 12.8)	9,808	77,587	12.6
2002	825	8,182	10.1	(9.4, 10.8)	9,202	76,929	12.0
2003	862	8,531	10.1	(9.4, 10.8)	8,624	79,328	10.9
2004	819	8,578	9.6	(8.9, 10.2)	8,068	79,265	10.2
2005	836	8,865	9.4	(8.8, 10.1)	8,174	80,302	10.2
2006	845	8,991	9.4	(8.8, 10.1)	8,712	84,498	10.3
2007	895	9,452	9.5	(8.9, 10.1)	8,818	87,041	10.1
2008	951	9,727	9.8	(9.2, 10.4)	9,174	89,429	10.3

<sup>1</sup> 95% Confidence Interval: A confidence interval is a range of values that has a specified probability of containing the rate. Narrow confidence intervals indicate that rate is precisely estimated, while wider confidence intervals indicate that the estimated rate is less precisely estimated.