

COVID-19 INTERVENTION TABLE BY TEST STATUS, EXPOSURE HISTORY, & CLINICAL STATUS

Current test status	Contact to COVID-19 in past 14d		≥1 current COVID-19 symptoms ¹		Disease control intervention
	Yes	No	Yes	No	
Positive	N/A ²	N/A ²	X		Isolate at home until: <ul style="list-style-type: none"> • Without fever for 24 hours³, AND • Other symptoms improving, AND • ≥10 days since onset of illness
	N/A ²	N/A ²		X	Isolate at home until 10 days after positive test specimen was collected
Negative	X		X		Isolate at home until: <ul style="list-style-type: none"> • Without fever for 24 hours³, AND • Other symptoms improving, AND • ≥10 days since onset of illness
	X			X	<ul style="list-style-type: none"> • Quarantine at home until 14 days after last exposure • Re-test promptly if symptoms emerge
		X	X		Isolate at home until: <ul style="list-style-type: none"> • Without fever for 24 hours³, AND • Other symptoms improving
		X		X	No isolation or quarantine necessary
Not done ⁴	X		X		Preferred: get tested and re-classify based on results Inferior alternative: isolate at home until: <ul style="list-style-type: none"> • Without fever for 24 hours³, AND • Other symptoms improving, AND • ≥10 days since onset of illness
	X			X	COVID-19 exposure without symptoms <ul style="list-style-type: none"> • Quarantine at home until 14 days after last exposure • Pursue testing if symptoms emerge
		X	X		Preferred: get tested and re-classify based on results Inferior alternative: isolate at home until <ul style="list-style-type: none"> • Without fever for 24 hours³, AND • Other symptoms improving, AND • ≥10 days since onset of illness
		X		X	<ul style="list-style-type: none"> • No isolation or quarantine necessary • No testing necessary

¹For current list of COVID-19 symptoms, go to <https://www.cdc.gov/coronavirus/2019-ncov/symptoms-testing/symptoms.html>

²Not applicable—does not impact disease control decision making.

³...without the use of fever reducing medications (e.g., acetaminophen, ibuprofen)

⁴COVID-19 testing is recommended (when available) for all contacts and all people with symptoms. If the personal health care provider is unwilling or unable to conduct testing, consider pursuing testing through SHD or another community-based testing venue (e.g., SeaMar, Community Health Centers, UW Medicine, other).